


Figure 1:



a: $p=0.024$ compared to group B control.

b: $p=0.008$ compared to group C restricted.

Figure 2:





Figure 3:



a: $p=0.019$ compared to group A fed animals.

Figure 4:




Figure 5:




Figure 6:






Figure 7:



a: $p=0.013$ compared to 17.5mM glucose.


b: $p=0.020$ compared to 25mM glucose.

Figure 8:




a: $p=0.044$ compared to 0.1nM insulin group.

Figure 9:




a: $p < 0.004$ compared to 0nM, 0.1nM, 1nM and 10nM insulin groups.

Figure 10:




a Gene expression significantly higher in B controls ($p=0.039$), C controls ($p=0.001$) when compared to A controls

Figure 11:




AGT 712 gene expression is positively correlated with post restriction glucose in control animals ($p=0.014$)

Figure 12:




AGT 712 gene expression is positively correlated with post restriction insulin in control animals ($p=0.016$)

Figure 13:



AGT 712 is Positively correlated with post restriction insulin in all animals ($p=0.042$)

Figure 14:



a: $p=0.039$

Figure 15:




Figure 16:



a: $p < 0.0001$

Figure 17:



a: p=0.005

Figure 18:




Figure 19:




Figure 20:




Figure 21:




Figure 22:




Figure 23:




Figure 24:





Figure 25:



a: $p=0.009$, Gene expression significantly lower in Group C animals compared to Group A and B animals.

b: $p=0.02$, Gene expression significantly lower in Group A fasted animals compared to Group A fed animals.

Figure 26:




Figure 27:






Figure 28:



a: Gene expression (mesenteric adipose tissue) significantly lower in A fasted animals when compared to A fed ($p=0.041$).

b: Gene expression (mesenteric adipose tissue) significantly lower in A fasted animals when compared to C fasted ($p=0.033$) animals.

Figure 29:



a: Gene expression (mesenteric adipose tissue) significantly lower in fasted animals when compared to fed animals ($p=0.014$).

Figure 30:






Figure 31:



a: Gene expression (3T3-L1 cells) significantly lower in 0nM insulin treated cells when compared to 0.1nM ($p=0.028$), 1nM ($p=0.046$), 10nM ($p<0.001$) and 100nM ($p=0.017$) insulin treated groups.

b: Gene expression (3T3-L1 cells) significantly higher in 10nM insulin treated cells when compared to 0nM ($p<0.001$), 0.1nM ($p=0.046$), 1nM ($p=0.028$), 1000nM and ($p=0.003$) insulin treated groups.


Figure 32:



a: Gene expression (3T3-L1 cells) significantly lower in 17.5mM glucose treated cells when compared to 0mM ($p=0.011$), 5mM ($p=0.01$) and 12.5mM ($p=0.011$) glucose treated groups.

b: Gene expression (3T3-L1 cells) significantly lower in 25mM glucose treated cells when compared to 0mM ($p=0.025$) and 5mM ($p=0.05$) glucose treated groups.

Figure 33:



a,b: Gene expression was significantly lower in A control animals compared to B control ($p=0.027$) and C control ($p<0.001$) animals.

c: $p=0.009$, gene expression was significantly higher in C control compared to C restricted animals.

Figure 34:




Figure 35:




Figure 36:




Figure 37:





Figure 38:



a: $p=0.023$, A control < C control

Figure 39:




Figure 40:




Figure 41:




Figure 42:




Figure 43:




Figure 44:






Figure 45:



a: A fed < B fed, C fed ($p=0.005, 0.007$)

b: A fasted < C fasted ($p=0.027$)

Figure 46:



a: Fed>Fasted ($p=0.038$)

Figure 47:




Figure 48:




Figure 49:






Figure 50:




a: B fed > C fed ($p=0.001$)

Figure 41:




a: Fed > Fasted ($p=0.007$)

Figure 52:



a: Fed > Fasted ($p=0.047$)

Figure 53:



a: $p < 0.05$, significant increase in the hypothalamus of B and C control groups, when compared with the A control group.

Figure 54:




Figure 55:




Figure 56:




Figure 57:




Figure 58:




Figure 59:




Figure 60:





Figure 61:



a: AGT-723 gene expression significantly higher ($p=0.005$) in C Control animals when compared to A Control.

b: AGT-723 gene expression significantly higher ($p=0.042$) in C Control animals when compared to B Control.

Figure 62:




Figure 63:




Figure 64:




Figure 65:




Figure 66:





Figure 67:



a: Hypothalamic *Psammomys obesus* AGT-723 gene expression (fed/fasted study) significantly higher ($p=0.032$) in B Fasted animals when compared to C Fasted animals.

Figure 68:






Figure 69:



a: Sprague Dawley rat hypothalamic AGT-723 gene expression (fed/fasted study) is significantly higher ($p=0.043$) in 48hr fasted animals when compared to 24hr fasted animals

Figure 70:



a: Mesenteric fat *Psammomys obesus* AGT-723 gene expression (fed/fasted study) significantly higher ($p=0.001$) in B Fed animals when compared to A Fed animals.
 b: Mesenteric fat *Psammomys obesus* AGT-723 gene expression (fed/fasted study) significantly higher ($p=0.005$) in B Fed animals when compared to B Fasted animals.
 c: Mesenteric fat *Psammomys obesus* AGT-723 gene expression (fed/fasted study) significantly higher ($p=0.022$) in B Fed animals when compared to C Fed animals.

Figure 71:




Figure 72:






Figure 73:



a,b: Gene expression was significantly lower in A controls when compared to B controls ($p=0.033$) and C controls ($p=0.004$)

Figure 74:






Figure 77:




Figure 78:




Figure 79:




Figure 80:




Figure 81:




Figure 82:






Figure 83:



a: $p=0.003$ compared to A fasted; $p=0.013$ compared to B fasted.

Figure 84:



a: $p<0.013$ compared to A fed, B fed and C fasted animals.

Figure 85:




Figure 86:




Figure 87:





Figure 88:



a: $p=0.018$ (A fed) compared to C fed.

b: $p=0.002$ (B fed) compared to C fed.

Figure 89:





Figure 90:



a: $p=0.022$, A control < C control

b: $p=0.040$, B control < C control

Figure 91:




Figure 92:




Figure 93:




Figure 94:




Figure 95:




Figure 96:




Figure 97:





Figure 98:




a: Gene expression significantly lower in A control animals compared to B control animals ($p=0.002$).


b: Gene expression significantly lower in A control animals compared to C control animals ($p=0.002$).

c: Gene expression significantly lower in A control animals compared to A restricted animals ($p=0.038$).


Figure 99:




Gene expression positively correlated with post restriction glucose ($p=0.020$) in control animals.

Figure 100:

Gene expression positively correlated with post restriction insulin ($p=0.034$) in control animals.

Figure 101:

Gene expression positively correlated with total activity ($p=0.045$) in control animals.

Figure 102:

Gene expression positively correlated with total fat ($p=0.017$) in control animals.


Figure 103:

Figure 104:

a: AGT-725 gene expression significantly lower in A controls when compared to C controls ($p = 0.012$)

b: gene expression significantly lower in C controls when compared to C energy restricted ($p=0.012$)




Figure 105:




Figure 106:




Figure 107: AGT-717 gene expression in all fed versus all fasted animals; *p=0.049 compared to fed animals.




Figure 108: Linear association between AGT-717 and glucose in red gastrocnemius muscle of P. obesus fasted for 24 hr.




Figure 109: AGT-717 gene expression in mesenteric fat of fed and 24 hr fasted P. obesus; * $p < 0.034$ compared to group A fed.




Figure 110: Linear associations of AGT-717 gene expression in mesenteric fat with body weight and insulin values in all animals.




Figure 111: AGT-717 gene expression in 3T3 cells treated with insulin for 24 hrs,
* $p < 0.01$ compared to 0 nM, 0.1 nM and 1 nM groups.





Figure 112: AGT-717 gene expression in 3T3 cells treated with glucose for 24 hr,
* $p < 0.05$ compared to 0 mM; # $p = 0.024$ compared to 12.5 mM.



**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.